

Calitar™

KEEP AWAY FROM MILK FEVER

FOR ANIMAL USE ONLY

|||||



KEEP AWAY FROM MILK FEVER KEEP AWAY FROM MILK FEVER KEEP AWAY FROM MILK FEVER KEEP AWAY FROM MILK FEVER KEEP AWAY FROM MILK FEVER

Ingredients:

Calcium chloride, calcium sulfate, xanthan gum, edible single double fatty acid ester.

Analytical content:

Item	Content, %
Calcium	≥22

Functions:

The effective prevention of clinical or subclinical hypocalcemia after calving and adjuvant therapy of clinical hypocalcemia, providing long-acting & sustained release of calcium supplement.

Using protocol:

Protocol 1	2nd parity dairy cows:	3rd parity dairy cows or above:
The prevention of clinical or subclinical hypocalcemia.	1 bolus immediately after calving.	1 st bolus immediately after calving. 2 nd bolus 12 hours after calving.
Protocol 2	For dairy cows with clinical signs of hypocalcemia:	
The adjuvant therapy of clinical hypocalcemia.	1 bolus 2-3 hours after intravenous injection treatment.	

Packing size:

200g net content per bolus.

When she delivers, Calitar™ protects always by her side.

How to use Calitar™ with QILU branded bolus applicator?

1. Place the bolus in the bolus applicator.
2. Gently guide the loaded QILU branded bolus applicator into the cow's mouth.
3. When the applicator is behind the tongue, squeeze the handle to release the bolus.
4. Wait a moment for the cow to swallow and then remove the applicator.

