

EnroQ

Enrofloxacin Injection

INDICATIONS

Treatment of infections of the respiratory tract caused by enrofloxacin susceptible strains of *Pasteurella multocida*, *Mycoplasma spp.* and *Actinobacillus pleuropneumoniae*. Treatment of infections of the urinary tract caused by enrofloxacin susceptible strains of *Escherichia coli*. Treatment of post-partum dysgalactiae syndrome, PDS (MMA syndrome) caused by enrofloxacin susceptible strains of *Escherichia coli* and *Klebsiella spp.* Treatment of infections of the alimentary tract caused by enrofloxacin susceptible strains of *Escherichia coli*. Treatment of septicaemia caused by enrofloxacin susceptible strains of *Escherichia coli*.

DOSAGE & ADMINISTRATION

2.5 mg of enrofloxacin/kg BW, once daily by intramuscular injection for 3 days. Alimentary tract infection, or septicaemia caused by *Escherichia coli*: 5 mg of enrofloxacin/kg BW, once daily by intramuscular injection for 3 days. In pigs, the injection should be made in the neck at the ear base. Not more than 3 mL should be administered at one intramuscular injection site.

CAUTIONS

- *Follow the prescription of veterinarian.
- *Not for human use.
- *Keep out of reach of children.

WITHDRAWAL TIME

Meat and offal 13 days

STORAGE CONDITIONS

Store in an airtight container, protected from light.

PACKING SIZE

10%

100 mL/vial; 50mL/vial

