

# QVECIN

## Cefovecin sodium for Injection



### Active Ingredients:

852 mg or 340mg cefovecin (as sodium salt) as a lyophilized cake.

### Indications:

Cefovecin sodium is indicated for the treatment of skin infections (secondary superficial pyoderma, abscesses, and wounds) in dogs caused by susceptible strains of *Staphylococcus intermedius* and *Streptococcus canis* (Group G). Cefovecin sodium is indicated for the treatment of skin infections (wounds and abscesses) in cats caused by susceptible strains of *Pasteurella multocida*. Cefovecin sodium is the first antibiotic for dogs and cats that ensures a course of treatment in a single injection. One dose of Cefovecin sodium provides up to 14 days of safe and effective antibiotic treatment for the most common skin infections. It is well-tolerated, and its safety has been demonstrated in juvenile and adult dogs and cats.

### Dosage & Administration:

**Dogs:** Cefovecin sodium should be administered as a single subcutaneous injection of 3.6 mg/lb (8 mg/kg) body weight. A second subcutaneous injection of 3.6 mg/lb (8 mg/kg) may be administered if response to therapy is not complete. The decision for a second injection for any individual dog should take into consideration such factors as progress toward clinical resolution, the susceptibility of the causative organisms, and the integrity of the dog's host-defense mechanisms. Therapeutic drug concentrations after the first injection are maintained for 7 days for *S. intermedius* infections and for 14 days for *S. canis* (Group G) infections. Maximum treatment should not exceed 2 injections.

**Cats:** Cefovecin sodium should be administered as a single, one-time subcutaneous injection at a dose of 3.6 mg/lb (8 mg/kg) body weight. After an injection of Cefovecin sodium, therapeutic concentrations are maintained for approximately 7 days for *Pasteurella multocida* infections.

### Adverse reaction:

**Dogs:** Lethargy, anorexia/decreased appetite, vomiting, diarrhea, blood in feces, dehydration, flatulence, increase borborygmi.

**Cats:** Vomiting, diarrhea, anorexia/decreased appetite, lethargy, hyper/acting strange, inappropriate urination.

### Cautions:

Not for use in humans. Keep it out of reach of children.

Dogs and cats allergic to  $\beta$ -lactam (penicillins and cephalosporins) group are not applicable.

For subcutaneous use in dogs and cats only.

### Specification:

852 mg/ vials, 340mg/ vials

### Storage Conditions:

Protect from light, 2° to 8° C.

Dry powder 24 months, reconstituted sterile solution 28 days.

If stored as recommended, solution color does not adversely affect potency.



FDA  
APPROVED



Vacuum  
Freeze-drying



Guarding  
Pets



Years of  
Concentration